BASF North Works
Phase II Sediment Investigation:

Preliminary Data Review & Proposed Next Steps

Helping Make Products Better™



Chicago, IL March 19, 2009



Presentation Outline

- Phase II Sediment Investigation Overview
- Core Collection & Processing
- Data Review & Analysis
 - Reference Areas
 - Along-Site Areas
- Preliminary Interpretations
- Next Steps & Schedule





Phase II Sediment Investigation Overview

- Objective: Evaluate sediment quality adjacent to North Works relative to upstream sediments
 - Collected & analyzed 26 upstream cores: 6 upstream of Mud Island, 5 near Mud Island, 2 in Ecorse River, 3 downstream of Ecorse River, 4 near Libra Marina, and 6 near Grassy Island
 - Collected & analyzed 30 cores from established along-site transects:
 1 from Northern Area (T-01 to T-03), 29 from Southern Area (T-19 to T-29)
 - Submitted samples from 3 upstream & 3 along-site cores for geochronological analysis



CORE COLLECTION & PROCESSING



Core Collection

- Vibracore sampling Oct. 14th-24th (2 cores in Libra Marina by hand Nov. 24th)
- Some upstream locations adjusted to increase sediment recovery; some abandoned and new locations substituted
- Upstream areas probed to locate potential depositional areas for core sampling

| Analytical Cores | Upstream¹ | Along-Site | Grassy Island |
|---------------------------|-----------------|-----------------|---------------|
| Proposed | 20 | 34 | 6 |
| Collected | 28 ² | 34 | 6 |
| Submitted for Analysis | 20 | 30 ³ | 6 |

Notes:

- 1) Upstream core count includes cores collected in Ecorse River and Libra Marina.
- 2) Collected upstream core count includes "A" and other locations with little sediment recovery that were subsequently relocated or replaced.
- 3) Locations T01-75; T02-25; T02-50; and T03-25 had very little sediment thickness and no samples were submitted for analysis. Probable depth previously observed was found to be composed of soft Native Clay versus sediments in these areas (Along-Site North).





Core Processing

- Cores temporarily stored and processed at Riverview facility
- Lab samples submitted per analyte list in the Work Plan







DATA REVIEW & ANALYSIS



Types of Data Collected During Field and Processing Activities

- Water Depth
- Probe Depth
- Recovery Depth
- Photographs
- Sediment Descriptions/Stratigraphy
- NAPL Observations
- Sediment Slurry pH
- Samples for Laboratory Analysis



U19 – Upstream of Mud Island NAPL Present at 2.9-3.2 feet



T27-50 (6-7.9 feet) Along-Site South Area



Upstream Probing & Core Locations





Upstream Core Field Data Summary

- Upstream pH Range: 6.9 to 10.5 *
 - Max upstream pH of 10.5 at U14 near entrance of Libra Marina
- Locations of greatest sediment deposition:
 - Upstream of Mud Island up to 8 ft
 - Downstream of Grassy Island up to 5 ft
- Sediment generally dark-brown or gray-brown silt, or fine sand over gray-brown silty clay
- Locations with very little to no sediment recovery:
 - U03, U08A, U09A (Near Mud Island)
 - U12, U06, U13A, U18 (Downstream of Ecorse River)
- NAPL observed in cores U04, U19, & U20 (upstream of Mud Island)

^{*} Note: All pH values in Standard Units (S.U.) and rounded to one decimal place.



Libra Marina Core Field Data Summary


- 1.5 to 4.1 ft of sediment recovery
- U14 pH range 9.5 to 10.5
- U15 pH range 8.1 to 9.5
- U16 & U17 located in Libra Marina
 - pH range from 7.2 to 9.0





Labadie Concrete (c.a. 1947-1973)

Libra Marina Area – Historical Aerial Photo circa 1937 (courtesy of MDEQ)






Libra Marina Core Examples

U17 - In Libra Marina



| U17 | |
|-----------------|--|
| Depth (feet) | Sediment Description |
| | Dark gray brown SILT, fine sand, trace |
| 0-1.3 | coarse sand, trace organics, odor |
| | Light to dark brown fine SAND, silt, trace |
| 1.3-2 | coarse sand, trace organics |
| | Dark brown to gray fine SAND, silt, trace |
| 2-2.9 | gravel, trace organics |




U16 - In Libra Marina



| U16 | |
|-----------------|---|
| Depth (feet) | Sediment Description |
| | Dark brown SILT, trace organics, trace fine sand, slight odor |
| | Brown fine SAND, some silt, trace organics, trace gravel |
| 2-2.6 | Gray to brown CLAY, trace fine sand, trace coarse sand, moderately soft, trace organics |


Upstream pH Data



U14 - Near Mouth of Libra Marina



| U14 | |
|-----------------|---|
| Depth (feet) | |
| (feet) | Sediment Description |
| | Dark gray inter-bedded fine SAND and SILT, trace medium to coarse sand, trace fine to |
| 0-1.5 | medium gravel, slight odor |






Along-Site Core Field Data Summary

- 11 Transects targeted
 - 3 near North area
 - 8 along South area
- 34 cores collected, 30 analyzed
- North Area: Transects 01, 02, & 03
 - 4 out of 5 cores collected had insufficient sediment thickness
 - Native Clay observed to be soft and probe-able
- South Area: Transect 19 through 29
 - Light colorations, crust fragments
 - Field testing indicates elevated pH







Transect 19





| T19-100 | |
|-----------------|--|
| Depth (feet) | Sediment Description |
| | Dark gray brown very loose SILT, little fine sand, trace fine gravel, trace NAPL blebs, slight odor |
| 0.4-3.7 | Light gray grading to white SILT, trace fine sand, odor |
| 3.7-3.9 | Light gray fine to coarse SAND, trace fine to medium gravel |







T23-50

0-0.5

Depth (feet) Sediment Description


odor

Dark gray SILT, trace fine sand,

Dark gray grading to light gray SILT, trace fine sand, odor




Transect 23



| T23-75 | |
|-----------------|--|
| Depth (feet) | Sediment Description |
| 0-2 | Blue-ish gray grading to light gray SILT, trace fine sand, odor |

















Transect 27














| | not sweensowel | |
|--|----------------|---------------|
| Ta | 7-100 | |
| | infac.let | |
| - Name to a real and | mportententen | andendandanks |
| A SO THE PROPERTY OF | 1 | La Sull |
| | | |
| 9 10 7 1 | Other Trans | |
| | - | |
| pH: 10.35 | 9.33 | 9.36 |

| 121-13 | |
|-----------------|--|
| Depth (feet) | Sediment Description |
| | Gray brown very loose SILT, |
| | trace fine sand, trace slag, odor, trace NAPL blebs |
| 3-3.8 | Dark gray SILT, trace fine sand, odor |
| 1 | Gray brown fine SAND, little light gray silt, odor |
| 5.4-6 | Dark gray brown SILT, trace fine sand, odor |

| T27-100 | |
|-----------------|--|
| Depth (feet) | Sediment Description |
| | Dark gray brown SILT, trace fine sand, trace organics (vegetation), odor |
| | Dark gray brown fine SAND, trace silt, odor |
| 2.4-3.3 | Light gray brown SILT, trace fine sand, odor |



pH Spatial Trends

- Upstream pH > 9 detected near Libra Marina area (unknown extent)
- Along-site pH > 9 limited to T19 to T29
- General increase in pH with depth
 - Max pH deeper than 1 ft in 76% of samples from T19 to T29
- General increase in pH values at stations further from shore
 - Max field pH of 11.9 recorded at location T23-100 (0.5-1.0 ft)
- Layer of light-colored material present at 10 core locations from T19 toT29 coinciding with elevated pH
 - Observed in 6 of 7 sample locations at 100 ft from shore



Considerations with Respect to pH

- No human exposure along-Site due to water depths, access restrictions
- Minimal surface water pH impacts expected due to River flows
- Ecosystem exposure primarily limited to benthos due to water depth¹
- Benthic toxicity throughout Detroit River documented by others²
 - Upstream stressors exist, pH an additional issue along-Site
- Exposure undetermined, unknown representativeness of data at depth
 - Natural sediments have deposited over the historical material
 - Mineral (CaCO₃) precipitate/crust should limit interface pH effects
 - Geochemical analysis indicates this should occur, site observations suggest this is present
- Representativeness of field pH measurements on sediment-water interface, and bio-available zone conditions need to be determined

Notes:

- 1) Average water depth in study area was 19.8 feet.
- 2) E.g., DREAMS 2000, Journal of Great Lakes Research 1996, Beak Consultants 1993.




Upstream v.s. Along-Site Data Comparisons

- Upstream reference area data compared to along-site samples to assess if discernable differences are present
 - Field pH readings
 - Laboratory analytical results
 - Considered effect of variable percentage of fine sediments due to river velocity variations (different depositional environments)
 - Aluminum, grain-size, and organic carbon normalization
- Graphical comparisons
 - Plots of summary statistics, concentration vs. position in river
- Calculation of Background Screening Levels (BSLs)
- Various statistical tests performed per the Work Plan





Sediment Textural Analysis





PRELIMINARY INTERPRETATIONS




Preliminary Upstream vs. Along-Site Data Comparison Results

- Data support hypothesis that along-site sediments are in a continuum of degraded sediment quality extending from upstream areas
 - Concentrations of many constituents (including mercury) are consistent with upstream sediments
- Discernable differences are apparent for some analytes, including pH
- Upstream data indicate source issues for a number of constituents
 - Benzene and 1,2-dichlorobenzene in Libra Marina
 - Metals upstream and in Ecorse River (e.g. vanadium and zinc)
 - PAHs and visual NAPL in upstream samples (U04, U19, U20)
 - Finer NAPL "blebs" in Along-Site cores also occur at upstream locations
 - Max Total PCBs¹ observed upstream: 46.4 mg/kg at U04 (0.5 to 1.0 ft)
 - pH near former Labadie Concrete (near current Libra Marina)



Example Boxplots Benzene, Chlorobenzene



Along Sita

Along Site (North)

Grassy Island

Libra Marina

† Result value is < 25th percentile - 3*IQR or > 75th percentile + 3*IQR

* Result_value is < 25th percentile - 1.5*IQR or > 75th percentile + 1.5*IQR

NA = value not applicable due to frequency of detection

ND = nondetec

IQR = interquartile range equals the 3rd quartile (75th percentile) - 1st quartile (25th percentile)

SD = standard deviation

ug/kg

ug/kg

ug/kg

Reporting limit is used for nondetects unless otherwise noted.

124 45.0

20 43.0

61 121 45.0

NA

94.0

56.0

140

NA

140

150

100.0

140

12.0

14.0

12.0

ND

ND

690

150

20.0

150

ND

ND

1,200

69.5

33.0

17.3

33.8

ND

ND

883

31.0

29.0

18.0

30.0

ND

ND

760

22.2

3.1

22.5

ND

ND

276

29.0

15.0

30.0

100.0

62.0

710

50.5

18.0

51.0

110

67.5

415

Values less than 10 are reported to 2 significant figures. Values greater than 10 are reported to 3 significant figures.

Notes

78.0 Along Site

Along Site (North)

Along Site (South)

19.5

78.0

135

74.0

760

55.3

- † Result value is < 25th percentile 3*IQR or > 75th percentile + 3*IQR
- * Result_value is < 25th percentile 1.5*IQR or > 75th percentile + 1.5*IQR
 - NA = value not applicable due to frequency of detection
 - ND = nondetect
 - IQR = interquartile range equals the 3rd quartile (75th percentile) 1st quartile (25th percentile)
 - SD = standard deviation

ug/kg

Reporting limit is used for nondetects unless otherwise noted.

119

116 45.0

20 43.0

19

59.0

94.0

56.0

140

70.0

150

100.0

10.0

ND

10.0

ND

ND

240

ND

65.0

ND

ND

2,400

ND

30.5

ND.

30.5

ND

ND.

1.210

ND

25.0

ND

25.0

ND

ND

1,000

Values less than 10 are reported to 2 significant figures. Values greater than 10 are reported to 3 significant figures.

58.0

61.0

58.0

100.0

62.0

72.0

65.0

74.0

110

67.5

190

76.0

84.0

68.8

84.5

135

74.0

1,000

84.3

17.5

ND

17.5

ND

ND


1,100

Helping Make
Products Better™

BASF

The Chemical Company

Example Boxplots Mercury and 4,4'-DDE



| | | | Sample Siz | e e | ND | tange | | | Detects | | | Perc | entiles (All | Data) | | | | Sample Siz | 00 | ND | Range | | | Detects | | | Perc | entiles (All | Data) |
|--------------------|-------|-----|------------|-------|-----|-------|------|-------|---------|--------|------|------|--------------|-------|--------------------|--------------|-----|------------|-------|-----|-------|------|-----|---------|--------|------|------|--------------|-------|
| Dataset | Units | NDs | Detects | Total | Min | Max | Min | Max | Mean | Median | SD | 25th | 50th | 75th | Dataset | ataset Units | NDs | Detects | Total | Min | Max | Min | Max | Mean | Median | .SD | 25th | 50th | 75th |
| Along Site | ug/kg | .0 | 130 | 130 | NA | NA | 14.0 | 3,200 | 907 | 735 | 742 | 260 | 735 | 1,300 | Along Site | ug/kg | 43 | 87 | 130 | 1.4 | 2.9 | 0.43 | 530 | 41.4 | 23.0 | 68.5 | 1.7 | 5.8 | 33.0 |
| Along Site (North) | ug/kg | 0 | 3 | 3 | NA | NA | 330 | 460 | 415 | 455 | 73.7 | 360 | 455 | 459 | Along Site (North) | ug/kg | 0 | 3 | 3 | NA | NA | 23.0 | 104 | 49.8 | 23.0 | 46.5 | 23.0 | 23.0 | 83.4 |
| Along Site (South) | ug/kg | 0 | 127 | 127 | NA | NA | 14.0 | 3,200 | 919 | 740 | 747 | 250 | 740 | 1,340 | Along Site (South) | ug/kg | -43 | 84 | 127 | 1.4 | 2.9 | 0.43 | 530 | 41.1 | 22.0 | 69.3 | 1.7 | 5.3 | 32.5 |
| Ecorse River | ug/kg | 0 | 7 | 7 | NA | NA | 59.0 | 1,050 | 527 | 510 | 335 | 310 | 510 | 765 | Ecorse River | ug/kg | 0 | 7 | 7 | NA | NA | 1.1 | 115 | 47.3 | 36.0 | 42.9 | 18.0 | 36.0 | 84.3 |
| Grassy Island | ug/kg | 0 | 20 | 20 | NA | NA | 44.0 | 1,300 | 387 | 185 | 370 | 140 | 185 | 655 | Grassy Island | ug/kg | 8 | 12 | 20. | 1.6 | 2.2 | 4.9 | 140 | 31.2 | 15.5 | 39.2 | 1.7 | 5.8 | 20.0 |
| Libra Manna | ug/kg | 0 | 6 | 6 | NA | NA | 140 | 760 | 418 | 425 | 205 | 300 | 425 | 460 | Libra Manna | ug/kg | 4 | 2 | 6 | 3.4 | 77.0 | 20.0 | 110 | 65.0 | 65.0 | 63.6 | 6.1 | 14.6 | 77.0 |
| Upstream | ug/kg | 0 | 69 | 69 | NA | NA | 41.0 | 3,200 | 1,180 | 1,000 | 777 | 610 | 1,000 | 1,730 | Upstream | ug/kg | 20 | 49 | 69. | 1.6 | 17.0 | 1.9 | 870 | 187 | 66.0 | 252 | 1.9 | 24.0 | 140 |

Notes


- † Result value is < 25th percentile 3*IQR or > 75th percentile + 3*IQR
- * Result_value is < 25th percentile 1.5*IQR or > 75th percentile + 1.5*IQR
 - NA = value not applicable due to frequency of detection
 - ND = nondetec
 - IQR = interquartile range equals the 3rd quartile (75th percentile) 1st quartile (25th percentile)
 - SD = standard deviation
 - Reporting limit is used for nondetects unless otherwise noted.
 - Values less than 10 are reported to 2 significant figures. Values greater than 10 are reported to 3 significant figures.


Notes:


- † Result value is < 25th percentile 3*IQR or > 75th percentile + 3*IQR
- * Result_value is < 25th percentile 1.5*IQR or > 75th percentile + 1.5*IQR
 - NA = value not applicable due to frequency of detection
 - ND = nondetect
 - IQR = interquartile range equals the 3rd quartile (75th percentile) 1st quartile (25th percentile)
 - SD = standard deviation
 - Reporting limit is used for nondetects unless otherwise noted.
 - Values less than 10 are reported to 2 significant figures. Values greater than 10 are reported to 3 significant figures.



Example Boxplots Vanadium and Zinc







| A A | RCAD | S |
|----------------|----------------|----------------|
| in Sautrycture | and the second | II, Donald St. |

VANADIUM

State Corporation, Wyandone, Mt

Figure C-16 ARCADIS

Sediment Box and Whisker Plot ZINC BASE Corporation, Wyandotte, MI Figure C-17

| Dataset | | Sample Size | | | ND Range | | Detects | | | | Percentiles (All Data) | | | | | Sample Size | | | ND Range | | Detects | | | | Percentiles (All Data) | | | | |
|--------------------|-------|-------------|-------|---------|----------|-----|---------|--------|--------|--------|------------------------|--------|--------|--------|--------------------|-------------|-----|---------|----------|------|---------|---------|----------|---------|------------------------|---------|---------|---------|---------|
| | Units | NDs | Delec | s Total | Min | Max | Min | Max | Mean | Median | SD | 25th | 50lb | 75th | Dataset | Units | NDs | Detects | Total | Min | Max | Min | Max | Mean | Median | SD | 25th | 50m | 75th |
| Along Site | ug/kg | 0 | 130 | 130 | NA: | NA | 7,700 | 27,500 | 16,500 | 17,000 | 4,340 | 13,000 | 17,000 | 19,800 | Along Sile | ug/kg | 0 | 130 | 130 | NA | NA | 12,300 | 732,000 | 263,000 | 264,000 | 183,000 | 100,000 | 264,000 | 389,000 |
| Along Site (North) | ug/kg | 0 | 3 | 3 | NA | NA | 12,900 | 15,150 | 14,000 | 14,000 | 1,130 | 13,000 | 14,000 | 14,900 | Along Site (North) | ug/kg | 0 | 3 | 3 | NA | NA | 304,000 | 376,000 | 336,000 | 327,000 | 36,800 | 310,000 | 327,000 | 364,000 |
| Along Site (South) | ug/kg | 0 | 127 | 127 | NA | NA | 7,700 | 27,500 | 16,600 | 17,100 | 4,370 | 13,000 | 17,100 | 19,800 | Along Site (South) | ug/kg | 0 | 127 | 127 | NA | NA | 12,300 | 732,000 | 262,000 | 238,000 | 185,000 | 98,000 | 238,000 | 392,000 |
| Ecorse River | ug/kg | 0 | 7 | 7 | NA | NA. | 22,700 | 33,600 | 28,500 | 27,900 | 4,130 | 25,000 | 27,900 | 32,400 | Ecorse River | ug/kg | 0 | 7 | 7 | NA . | NA | 67,500 | 741,000 | 365,000 | 393,000 | 208,000 | 240,000 | 393,000 | 406,000 |
| Grassy Island | ug/kg | 0 | 20 | 20 | NA | NA. | 10,100 | 33,100 | 18,400 | 17,400 | 7,110 | 14,000 | 17,400 | 21,800 | Grassy Island | ug/kg | 0 | 20. | 20 | NA | NA | 48,900 | 299,000 | 126,000 | 88,900 | 79,200 | 69,000 | 88,900 | 171,000 |
| Libra Marma | ug/kg | 0 | 6 | 6 | NA. | NA. | 20,100 | 41,400 | 28,900 | 27,800 | 7.020 | 26,000 | 27,800 | 30,100 | Libra Manna | ug/kg | 0 | 6 | 6 | NA | NA | 129,000 | 602,000 | 369,000 | 367,000 | 163,000 | 290,000 | 367,000 | 458,000 |
| Upstream | ua/ka | 0 | 69 | 69 | NA | NA. | 8,000 | 36,700 | 22,800 | 22,400 | 6,490 | 20,000 | 22,400 | 26,900 | Upstream | ug/kg | 0 | 69 | 69 | NA | NA | 38,800 | 1.17E+06 | 540,000 | 543,000 | 230.000 | 450,000 | 543,000 | 690,000 |

Notes

- \uparrow Result value is < 25th percentile 3*IQR or > 75th percentile + 3*IQR
- * Result_value is < 25th percentile 1.5*IQR or > 75th percentile + 1.5*IQR
 - NA = value not applicable due to frequency of detection
 - ND = nondetec
 - IQR = interquartile range equals the 3rd quartile (75th percentile) 1st quartile (25th percentile)
 - D = standard deviation
 - Reporting limit is used for nondetects unless otherwise noted.
 - Values less than 10 are reported to 2 significant figures. Values greater than 10 are reported to 3 significant figures.

Notes

- † Result value is < 25th percentile 3*IQR or > 75th percentile + 3*IQR
- * Result_value is < 25th percentile 1.5*IQR or > 75th percentile + 1.5*IQR
 - NA = value not applicable due to frequency of detection
 - ND = nondetect
 - IQR = interquartile range equals the 3rd quartile (75th percentile) 1st quartile (25th percentile)
 - SD = standard deviation
 - Reporting limit is used for nondetects unless otherwise noted.

Values less than 10 are reported to 2 significant figures. Values greater than 10 are reported to 3 significant figures



Preliminary Results of Upstream Vs. Along-Site Statistical Comparisons

Inorganics


- Normalized to Aluminum (Al) using bivariate plots for 17 inorganics
 - Preliminary BSLs established based on 95% Prediction Interval (P.I.) for comparison to Along-Site values
- Preliminary BSLs not exceeded for mercury, antimony, barium, cadmium, chromium, cobalt, copper, nickel, silver, vanadium, zinc
 - Consistent with Conceptual Site Model description of continuum of impacts from upstream sources
- Preliminary BSLs exceeded by varying percent of total sample counts for arsenic, beryllium, lead, selenium, and thallium and cyanide
 - Suggests potential localized, incremental impact for some constituents
- Addition evaluation (e.g. hypothesis testing) is currently being concluded


Organics

- VOCs, SVOCs, PAHs, Total PCBs, 4,4'-DDE
- Evaluation is currently being concluded
- Normalized to TOC



Example Al Normalization and Prediction Interval Plots







NEXT STEPS & SCHEDULE



Next Steps

- Complete Draft Phase II Sediment Investigation Report for Agency Review
- Develop plan outline to address data needs, determine exposure conditions in Study Area
- Provide plan outline to USEPA, MDEQ
 - Conference call with Agencies to discuss approaches
 - Submit Draft Work Plan for Agency review
- Explore and evaluate options for addressing sediment concerns
- Collect supplemental data needed to close data gaps
- Meet with USEPA and MDEQ to review additional data, considerations on options, and path forward



Supplemental Data Collection

- Core sampling for field measurement of pH and visual observation
 - Delineate elevated pH around Libra Marina
 - Delineate elevated pH in Along-Site Area
 - Determine depth to top of clay
 - Collect step-out cores beyond 100 ft for additional analysis
- Near-bottom surface water pH measurements
- Sediment-water interface pH measurements
 - Assess bioavailable zone conditions
- Evaluate Surface Sediment Geochemistry
- Sediment Profile Imaging (SPI) survey
- Multi-beam sonar bathymetric survey



Example SPI photograph of benthic zone after disposal of drilling mud (light sand material over dark drilling mud)



Estimated Schedule

